

ORAL MICROBIOLOGY

Laboratory and clinical comparison of preservation media and transport conditions for survival of *Actinobacillus actinomycetemcomitans*

R. PICCOLOMINI, G. CATAMO, G. DI BONAVENTURA, C. PICCIANI
and MICHELE PAOLANTONIO*

Department of Biomedical Sciences, Clinical Microbiology Laboratory and *School of Dentistry; 'G. D'Annunzio' University, Via dei Vestini, 31 I-66100-Chieti, Italy

The capacity of clinical isolates and type strains of *Actinobacillus actinomycetemcomitans* to survive in a new transport medium (AaTM), phosphate-buffered saline (PBS) and Ringer's solution (RS) was evaluated. The effects of exposure to air, transportation time and temperature on viability were also studied. In addition, the culture of *A. actinomycetemcomitans* from subgingival plaque of patients with different forms of periodontitis was quantified. The results following storage in AaTM, PBS and RS showed that *A. actinomycetemcomitans* survived better in AaTM than in PBS or RS when transportation times exceeded 20–22 h, and that survival was enhanced by storage at below 12°C. Serotype b strains of *A. actinomycetemcomitans* were able to survive better than either serotype a or c. In the clinical study the optimal transportation conditions for subgingival plaque containing *A. actinomycetemcomitans* were AaTM at a temperature of 8°C for 24 h under anaerobic conditions. These conditions resulted in a high survival and isolation rate for *A. actinomycetemcomitans* without inhibition of the other periodontopathic bacteria isolated from deep periodontal pockets. These findings have practical implications for future multicentre clinical trials in which the transportation of oral specimens over relatively long distances and at different ambient temperatures during various periods of the year are required.

Introduction

Actinobacillus actinomycetemcomitans has been strongly implicated in the pathogenesis of juvenile and adult periodontitis as well as endocarditis [1]. Therefore, microbiological monitoring of patients with periodontal diseases has been employed in diagnosis, choice of therapy, treatment control and risk assessment [2, 3]. Even if other methods, such as molecular diagnosis, are available as an alternative procedure in periodontal infections, the culture of *A. actinomycetemcomitans* is the only diagnostic method that allows assessment of the susceptibility of this organism to antimicrobial agents.

For maximal recovery of *A. actinomycetemcomitans* from clinical specimens, it is desirable to culture subgingival samples without delay. As this is not always possible because of the distance between non-

institutional dental centres and clinical microbiology laboratories, a few studies have evaluated the various conditions required to transport oral plaque over relatively long periods of time. The importance of certain factors, such as metabolic requirements, pH, salt concentration and the effects of freeze-thawing on the growth and viability of *A. actinomycetemcomitans* has been reported [4]. It has also been observed that a delay of 1–2 days in the transportation of plaque samples in Ringer's solution from non-institutional dental centres yielded poor rates of isolation of *A. actinomycetemcomitans*. Furthermore, the survival of *A. actinomycetemcomitans* was influenced by the duration of exposure to air before culture [5].

Currently, various transport media are available for the preservation of oral samples containing *A. actinomycetemcomitans* and other periodontopathic bacteria under micro-aerophilic conditions [6–8], but none is ideal, mainly because of their inability to stabilise bacterial cells in the samples without allowing them to proliferate. The conditions required for the transportation of oral specimens before culture influence a

Received 31 July 1997; revised version accepted 16 Dec. 1997.

Corresponding author: Dr R. Piccolomini.

crucial phase in the microbiological assessment of periodontal infections. Consequently, it is essential to employ a transport medium that can maintain the complex subgingival flora both qualitatively and quantitatively.

The new medium was designed to improve the anaerobic transport of subgingival plaque micro-organisms, especially important periodontal pathogens such as *A. actinomycetemcomitans*, and was compared with phosphate-buffered saline (PBS) and Ringer's solution (RS) under various experimental conditions. Subsequently, the recovery of *A. actinomycetemcomitans* and other oral micro-organisms from the subgingival plaque of patients with periodontal diseases following transportation in these three media was compared.

Materials and methods

Laboratory study

Micro-organisms. Three type cultures of *A. actinomycetemcomitans* (ATCC 29522, ATCC 29523 and NCTC 9710) obtained from the American Type Culture Collection (Rockville, MD, USA) and from the National Collection of Type Cultures (London) and 10 wild strains were tested. The latter were randomly chosen fresh strains isolated from periodontal pockets of patients with localised juvenile periodontitis who were being treated in the School of Dentistry at the 'G. D'Annunzio' University. Clinical isolates were designated *A. actinomycetemcomitans* if they possessed the following characteristics: gram-negative, facultatively anaerobic short rods; NAD, haemin, oxidase, urease and tryptophanase negative, catalase positive and produced acid from glucose but not from lactose or sucrose [9, 10]. In addition, the small, convex, adherent colonies on TSBV-agar plates [11] exhibited typical morphology when examined in oblique or transmitted illumination with $\times 10$ –80 magnification in a Leica zoom stereomicroscope.

The serotype of *A. actinomycetemcomitans* isolates was confirmed by serology [12]. The methods for raising antisera against *A. actinomycetemcomitans* strain ATCC 29523 (serotype a), strain ATCC 29522 (serotype b) and strain NCTC 9710 (serotype c) were those described by Asikainen *et al.* [13]. Bacterial cells were grown in 5 ml of Todd-Hewitt Broth (Oxoid, Garbagnate Milanese, Milan, Italy) supplemented with yeast extract 1.2% w/v at 36°C for 3 days in an atmosphere containing CO₂ 5% in air. Antigenic extracts were prepared by autoclaving the *A. actinomycetemcomitans* cells at 121°C for 15 min [14]. Immunodiffusion was performed in Noble 1.2% agar by the Ouchterlony double-diffusion method [15].

For long-term storage, subcultures originating from one colony were frozen in Trypticase Soy Broth

(Oxoid) supplemented with yeast extract (Oxoid) 0.5% w/v and NaHCO₃ 0.05% w/v containing dimethylsulphoxide 10%.

Transport media. *A. actinomycetemcomitans* transport medium (AaTM) was composed of bacteriological gelatin (LP 008B; Oxoid) 80 g/L, and re-distilled water 500 ml, to which were added the following compounds, obtained from Sigma-Aldrich (Milan, Italy): thioglycollic acid-sodium salt 1.5 g/L, sodium phosphate 1.1 g/L, NaCl 5.1 g/L, KCl 0.30 g/L, calcium chloride dihydrate 0.15 g/L, methylene blue 3 mg/L and re-distilled water 490 ml. This mixture was boiled on a hotplate while bubbling oxygen-free CO₂ gas into the molten transport medium through a Pasteur pipette, for about 4 min, which was then replaced by oxygen-free N₂ for a further 2 min. When the methylene blue indicator had become colourless, L-cysteine (12 mg) was added, and dissolved by gentle heating in 10 ml of distilled water. Finally, the pH of the medium was adjusted to 7.5 with boiled then cooled 0.1 N NaOH:0.1 N KOH (1:1). The medium was transferred immediately to an anaerobic chamber (Don Whitley Scientific; International PBI SpA, Milan, Italy) and dispensed into sterile pre-reduced 8-ml vials. After tight capping, the vials were autoclaved at 115°C for 20 min. Vials with an azure or blue colouration were discarded.

PBS was prepared as described by Dulbecco and Vogt in Sambrook *et al.* [16] and Ringer's solution (RS; Oxoid) was prepared according to the manufacturer's instructions. PBS (pH 7.3) and RS (pH 7.0) were dispersed in 8-ml screw-capped vials and oxygen-free N₂ and CO₂ gas was added to the space between the medium surface and the vial top as described above for AaTM. All vials were used within 2 weeks of preparation.

Exposure to air. Fresh strains of *A. actinomycetemcomitans* were cultivated in anaerobic jars (95% N₂, 5% CO₂) for 72–96 h on TSBV agar at 37°C and harvested from plates by scraping, then suspended in 0.1 M phosphate buffer (pH 6.8) to a density of *c.* 10⁸ cells/ml, corresponding to a McFarland opacity standard of 0.5 [17]. The time required for the preparation of the bacterial suspensions was no longer than 3 min. To evaluate the effect of exposure to air on survival, 30-ml samples from the suspension were dispensed in opened tissue-culture roller bottles (International PBI) then exposed to air and agitated in a roller culture apparatus (W. Pabisch SpA, Milan, Italy) at room temperature (18°C \pm 1°C) in a sterile laminar flow cabinet (Gruppo Flow SpA, Opera-Milan, Italy) for 15, 30 and 60 min. After each period of air exposure, 100 μ l were removed to measure survival rate, and following 10-fold serial dilutions in 0.1 M phosphate buffer, samples were inoculated on to three TSBV-agar plates which were incubated in anaerobic jars (95% N₂, 5% CO₂) at 37°C for 5 days. Culture

plates were selected with *c.* 300 colonies/plate, and after counting the numbers present were then expressed as cfu/ml.

Holding time and temperature. *A. actinomycetemcomitans* strains were cultivated as described above for 72–96 h on TSBV agar at 37°C then suspended in 0.1 M phosphate buffer (pH 6.8) to a density corresponding to *c.* 10⁸ cells/ml. A standard inoculum (100 µl) of the suspensions prepared for each of the 13 test strains was immediately absorbed on to 12 fine paper points (Dentsply DeTrey, Rome, Italy) which were transferred to four vials of each medium: (i) *Aa*TM, (ii) PBS or (iii) RS. Vials were stored for various periods of time (8, 16 and 24 h) and at different temperatures (5, 12, 20 and 28°C) before subsequent culture and recovery. All *Aa*TM vials were incubated at 28°C for 20 min before the contents were mixed in a vortex mixer for 30 s. Samples (100 µl) from the undiluted transport medium and from a 100-fold dilution in 0.1 M phosphate buffer were spread on TSBV-agar for incubation and subsequent estimates of viability.

Clinical study

Patients and bacteriological procedures. Seventy patients with different forms of early-onset and adult periodontitis were included in this part of the study. None of the subjects had a history of serious illness nor had they received antibiotics for at least 4 months before the microbiological investigation. The most diseased site (pocket depth of ≥6 mm) in each quadrant was selected for bacteriological sampling, which was performed immediately after recording plaque indices and before the assessment of the gingival bleeding index [18]. After removal of supragingival plaque by a scaler, the gingival surface was dried by a gentle flow of sterile, oxygen-free CO₂ gas. Plaque samples were collected by the insertion of three sterile fine paper points into the periodontal pocket. These were left *in situ* for 15 s then removed at the same time; each was immediately transferred into a vial containing 1.0 ml of one of the three test transport media. The paper points from each quadrant were added to the same vial to produce a pooled 'mouth' sample for each patient. Plaque samples were maintained at a temperature of +8°C (±1°C) in a portable refrigerator (Penguin TR825H; PBI International, Milan, Italy) and transported to the Clinical Microbiology Laboratory of the 'G. D'Annunzio' University of Chieti where they were processed 24 h after collection.

Samples in the *Aa*TM vials were warmed to 28°C for 20 min before processing. All samples were mixed for 30 s in a vortex mixer at the maximal setting then diluted 1 in 10 in 0.1 M phosphate buffer. Samples (100 µl) of the appropriate dilution of each pooled plaque sample were inoculated in triplicate on to freshly prepared TSBV-agar plates, Columbia Blood

Agar Base (CBA; Oxoid) enriched with non-spore-forming anaerobic bacteria supplement (SR 107B + Tween 80; Oxoid) and Wilkins-Chalgren Anaerobe Blood Agar (WCAB; Oxoid) enriched with gram-negative anaerobic bacteria supplement (SR 108B; Oxoid) for total anaerobic colony counts. The plates were then cultured at 37°C in a CO₂ 5% incubator (TSBV), aerobically (CBA), and in an anaerobic chamber (WCAB) for 5 days, then examined for the presence and enumeration of *A. actinomycetemcomitans* and the presumptive identification of aerobic and anaerobic micro-organisms from plaque. Plates containing 30–300 cfu were selected for quantitative estimation of bacterial growth.

A. actinomycetemcomitans on TSBV-agar was identified by the criteria described previously. Initial differentiation of the black-pigmented *Porphyromonas* and *Prevotella* spp. was based on cell and colony morphology, pigment production, fluorescence under long-wave UV light [19], and susceptibility or resistance to special-potency antibiotic disks (vancomycin 5 µg; kanamycin 1000 µg; colistin 10 µg; Oxoid) [20]. Colonies of gram-negative filamentous, spindle-shaped and non-motile rods were presumptively classified as *Fusobacterium* spp. and were easily differentiated from *Capnocytophaga* spp. because of the micro-aerophilic characteristic of the latter. Gram-positive *Peptostreptococcus* spp. were preliminarily identified on the basis of colony morphology on the WCAB plates as small white colonies with haemolysis. A definitive identification of all representative isolates was then obtained by subculturing on to Brucella Blood Agar (Oxoid) followed by inoculation of purified cultures into a commercially packaged automated system (Vitek; bioMérieux Italia SpA, Rome, Italy).

The proportion of *A. actinomycetemcomitans* and the aerobic microflora present in the pooled plaque samples were determined by dividing the number of typical colonies on TSBV-agar and the total cfu on CBA, respectively, by the total anaerobic colony counts on the WCAB plates; the results being reported as a percentage of the total anaerobic count.

Results

Laboratory study

Influence of exposure to air. The effect of exposure to air on the viability of *A. actinomycetemcomitans* is presented in Table 1. The type strains tested were more aerotolerant than fresh clinical isolates and after exposure to air for 30 min, both the size and number of colonies recovered was greater than that achieved with any of the 10 clinical isolates (seven showed moderate growth and three scanty growth). When exposure to air was extended to 60 min, the majority of the strains tested (84.6%) could not be subsequently

Table 1. Recovery of three type strains and 10 clinical isolates of *A. actinomycetemcomitans* in 0.1 M phosphate buffer after various holding times with exposure to air at room temperature ($18^{\circ}\text{C} \pm 1^{\circ}\text{C}$)

Strain (serotype)	Recovery after holding for		
	15 min	30 min	60 min
ATCC 29522 (b)	3+	2+	±
ATCC 29523 (a)	3+	2+	—
NCTC 9710 (c)	3+	2+	—
UdA 06 (a)	2+	±	—
UdA 07 (b)	3+	+	—
UdA 12 (c)	3+	+	—
UdA 14 (a)	2+	±	—
UdA 15 (b)	3+	+	—
UdA 18 (b)	3+	+	—
UdA 23 (b)	3+	+	±
UdA 25 (c)	2+	+	—
UdA 37 (b)	3+	+	—
UdA 43 (a)	2+	±	—

An inoculum of 10^6 cfu/ml was used and recovery was graded as follows: 3+, abundant ($>10^5$ cfu/ml); 2+, good (10^5 – 10^4 cfu/ml); +, moderate (10^3 – 10^2 cfu/ml); ±, scanty ($<10^2$ cfu/ml); —, no growth.

recovered and two strains, a type strain and a fresh clinical isolate both belonging to serotype b, grew poorly.

Influence of medium composition, holding time and temperature. The recovery of *A. actinomycetemcomitans* from AaTM was superior to that obtained from other transport media (Table 2). After 8 h at holding temperatures of 5, 12 and 20°C , all 13 strains were recovered from AaTM, compared with six from PBS and none from RS. The number of clinical strains of *A. actinomycetemcomitans* isolated from AaTM was 10^2 – 10^4 times greater than the numbers isolated from either PBS or RS after 16 h at 12 and 20°C . When the temperature was increased to 28°C , the majority of *A. actinomycetemcomitans* strains tested failed to grow in either PBS or RS; conversely, the serotype b strains shown moderate growth in AaTM (data not shown). No strain of *A. actinomycetemcomitans* survived incubation in PBS or RS at 20°C for 24 h, whereas five of the

Table 2. Recovery of 13 strains of *A. actinomycetemcomitans* from transport medium (AaTM), PBS and Ringer's solution (RS) after exposure to different temperatures for 8, 16 and 24 h

Strain (serotype)	Exposure (h)	Recovery after holding at								
		5°C			12°C			20°C		
		AaTM	PBS	RS	AaTM	PBS	RS	AaTM	PBS	RS
ATCC 29522 (b)	8	3+	3+	3+	3+	3+	±	3+	+	—
	16	3+	3+	+	3+	2+	±	2+	+	—
	24	3+	+	±	2+	+	—	+	±	—
ATCC 29523 (a)	8	3+	3+	2+	3+	3+	±	2+	—	—
	16	3+	2+	±	3+	+	—	+	—	—
	24	2+	+	—	+	—	—	—	—	—
NCTC 9710 (c)	8	3+	2+	+	3+	2+	±	2+	—	—
	16	2+	2+	±	2+	+	—	+	—	—
	24	2+	±	—	±	—	—	—	—	—
UdA 06 (a)	8	3+	3+	2+	3+	2+	±	2+	—	—
	16	2+	2+	±	2+	±	—	±	—	—
	24	+	±	—	±	—	—	—	—	—
UdA 07 (b)	8	3+	3+	3+	3+	2+	±	3+	+	—
	16	3+	3+	+	3+	+	±	2+	±	—
	24	3+	2+	—	+	—	—	±	—	—
UdA 12 (c)	8	3+	2+	+	2+	+	—	+	—	—
	16	2+	2+	—	+	±	—	±	—	—
	24	+	±	—	±	—	—	—	—	—
UdA 14 (a)	8	3+	2+	2+	2+	+	—	2+	—	—
	16	3+	2+	—	2+	±	—	+	—	—
	24	2+	±	—	+	—	—	—	—	—
UdA 15 (b)	8	3+	3+	2+	3+	2+	+	3+	+	—
	16	3+	3+	+	3+	+	±	+	±	—
	24	3+	2+	±	+	±	—	—	—	—
UdA 18 (b)	8	3+	3+	2+	3+	2+	+	3+	±	—
	16	3+	3+	+	3+	+	±	2+	±	—
	24	3+	2+	±	2+	±	—	+	—	—
UdA 23 (b)	8	3+	3+	3+	3+	2+	+	3+	±	—
	16	3+	3+	+	3+	+	±	2+	—	—
	24	3+	+	—	2+	—	—	+	—	—
UdA 25 (c)	8	3+	2+	2+	2+	+	—	+	—	—
	16	2+	+	—	+	±	—	—	—	—
	24	+	—	—	±	—	—	—	—	—
UdA 37 (b)	8	3+	3+	2+	3+	3+	+	3+	+	—
	16	3+	3+	±	3+	+	—	+	—	—
	24	3+	+	—	2+	—	—	+	—	—
UdA 43 (a)	8	3+	2+	2+	2+	2+	—	2+	—	—
	16	2+	+	—	2+	±	—	±	—	—
	24	2+	—	—	+	—	—	—	—	—

See footnote to Table 1.

13 strains were recovered from *Aa*TM. A clear correlation between temperature, time and recovery was demonstrated; and providing that the holding period was <8 h, the re-isolation of all 13 strains from *Aa*TM was possible at temperatures of up to 20°C. If the holding time was >8 h, the influence of temperature became critical, as only four of the 10 clinical isolates could be recovered in small numbers after 16 h at 28°C. After storage for 24 h, all strains of *A. actinomycetemcomitans* tested survived when they were held in *Aa*TM at 12°C, but only three serotype b strains were recovered from PBS and none from RS.

Clinical study

The total viable anaerobic count of plaque samples from deep periodontal pockets on WCAB plates was significantly higher in samples held in *Aa*TM than in PBS or in RS ($p < 0.01$; Wilcoxon rank sum test) (Table 3). The mean percentage of *A. actinomycetemcomitans* of the total viable plaque counts was 10.3% when samples were stored for 24 h at 8°C in *Aa*TM. This was consistently higher than those exhibited by plaque samples stored under the same conditions in PBS (2.1%, $p < 0.05$) or in RS (1.9%, $p < 0.01$).

Other potential pathogens (*F. nucleatum*, *P. gingivalis*, *Prev. intermedia* and *Prev. melaninogenica*) were also isolated in significantly higher numbers ($p < 0.05$) from 70 subgingival plaque samples when stored in *Aa*TM for 24 h at 8°C, compared with PBS or RS (data not shown). However, *Peptostreptococcus* spp., *Capnocytophaga* spp., *Candida albicans* and certain coliforms were more frequently isolated from PBS or RS than from *Aa*TM.

Discussion

A. actinomycetemcomitans is a fastidious periodontopathic micro-organism and for optimal isolation from subgingival plaque it is essential to culture the sample as soon as possible. However, delay is usually unavoidable and subgingival plaque is placed temporarily in transport medium. It is likely that a serious limitation to the efficient isolation of *A. actinomycetemcomitans* from oral specimens is the cumulative effect of inappropriate transport conditions, such as the

duration of exposure to air, the delay between sampling and culture, and incorrect transport temperature. This is supported by studies which show that *A. actinomycetemcomitans* is able to survive for up to 20 min in PBS at 22°C, but survives better at low temperatures [4]. Pre-reduced anaerobically sterilised Ringer's solution has been employed for transporting subgingival plaque [21], but <1% of *A. actinomycetemcomitans* was detected from the total culturable microbiota. It appeared that atmospheric levels of oxygen were toxic for *A. actinomycetemcomitans*, especially in the case of serotypes a and c.

In the present study, oxygen was excluded from all three transport media at different stages, but *Aa*TM proved to be most effective, which may be due in part to the presence of gelatin (8%) in *Aa*TM. This was liquified by heating samples to 28–30°C followed by vortex mixing before inoculation. *Aa*TM allows plaque micro-organisms to survive better than either PBS or RS at holding times that exceed 20–22 h, and this survival is enhanced at storage temperatures below 12°C. These results indicate that the recovery rates of *A. actinomycetemcomitans* were significantly higher when *Aa*TM was used. All strains of *A. actinomycetemcomitans* stored in *Aa*TM survived for as long as 24 h at 8°C, permitting the successful recovery of the micro-organism without inhibition of the other periodontopathic bacteria (such as *P. gingivalis*, *Prev. melaninogenica*, *Prev. intermedia* and *F. nucleatum*) isolated from deep periodontal pockets.

Xylose positive serotype b strains of *A. actinomycetemcomitans* were able to survive in *Aa*TM better than serotypes a and c in the laboratory study. The reason for this result is unknown, but may be due to the presence of a physicochemical complex present only in serotype b cells that protects against exposure to air, holding time and temperature. This could also explain in part why serotype b strains of *A. actinomycetemcomitans* have been isolated more commonly from subgingival plaque of patients with localised juvenile periodontitis than either serotype a or c [22].

As only one optimal, but expensive and complex transport medium suitable for mailing of *A. actino-*

Table 3. Total viable count (anaerobic) and total aerobic count and mean percentage of *A. actinomycetemcomitans* and aerobic microflora of total viable count in 70 subgingival plaque samples held for 24 h in *Aa*TM, PBS and RS at 8°C ($\pm 1.0^\circ\text{C}$) before culture

Transport medium	Mean (range) total count cfu $\times 10^6$		Mean (range) percent of total anaerobic count	
	anaerobic	aerobic	<i>A. actinomycetemcomitans</i>	aerobic flora
<i>Aa</i> TM	18.4 (1.46–96.4)	7.6 (0.49–28.6)	10.3 (1.6–32.4)	41.3 (20.6–79.3)
PBS	6.2 (0.051–21.2)	1.3 (0.035–5.6)	2.1 (0.9–7.6)	20.9 (19.3–81.2)
RS	4.6 (0.057–19.8)	0.92 (0.030–6.3)	1.9 (0.7–3.8)	20 (18.6–80.6)

mycetemcomitans has been developed previously [23], AaTM might represent an alternative for this purpose, although the need to maintain samples between 8 and 12°C and limit the transportation time to <24 h makes it less convenient to employ. Also, some species did not survive well in AaTM (*Peptostreptococcus* spp., *Capnocytophaga* spp. and *Candida* spp.). Therefore, a study comparing the medium described by Dahlen *et al.* [23] and AaTM is required so that the best medium and transportation conditions for conveying plaque samples over long distances can be identified.

We thank Carmelita Della Pelle, Simonetta D'Ercole and Stefania Lupo for their technical assistance. This work was supported by grants from the Ministero dell'Università e Ricerca Scientifica e Tecnologica (MURST, 'ex-*quote* 40% e 60%').

References

- Schack SH, Smith PW, Penn RG, Rapoport JM. Endocarditis caused by *Actinobacillus actinomycetemcomitans*. *J Clin Microbiol* 1984; **20**: 579–581.
- Listgarten MA. A rationale for monitoring the periodontal microbiota after periodontal treatment. *J Periodontol* 1988; **59**: 439–444.
- Maiden MFJ, Carman RJ, Curtis MA *et al.* Detection of high-risk groups and individuals for periodontal diseases: laboratory markers based on the microbiological analysis of subgingival plaque. *J Clin Periodontol* 1990; **17**: 1–13.
- Sreenivasan PK, Meyer DH, Fives-Taylor PM. Factors influencing the growth and viability of *Actinobacillus actinomycetemcomitans*. *Oral Microbiol Immunol* 1993; **8**: 361–369.
- Paolantonio M, di Girolamo G, Pedrazzoli V *et al.* Occurrence of *Actinobacillus actinomycetemcomitans* in patients wearing orthodontic appliances. A cross-sectional study. *J Clin Periodontol* 1996; **23**: 112–118.
- Helstad AG, Kimball JL, Maki DG. Recovery of anaerobic, facultative, and aerobic bacteria from clinical specimens in three anaerobic transport systems. *J Clin Microbiol* 1977; **5**: 564–569.
- Hill GB. Effects of storage in an anaerobic transport system on bacteria in known polymicrobial mixtures and in clinical specimens. *J Clin Microbiol* 1978; **8**: 680–688.
- Spiegel CA, Minah GE, Krywolap GN. Improved procedure for transport of dental plaque samples and other clinical specimens containing anaerobic bacteria. *J Clin Microbiol* 1979; **9**: 637–639.
- Slots J. Salient biochemical characters of *Actinobacillus actinomycetemcomitans*. *Arch Microbiol* 1982; **131**: 60–67.
- Slots J. Rapid identification of important periodontal microorganisms by cultivation. *Oral Microbiol Immunol* 1986; **1**: 48–57.
- Slots J. Selective medium for isolation of *Actinobacillus actinomycetemcomitans*. *J Clin Microbiol* 1982; **15**: 606–609.
- Amano K, Nishihara T, Shibuya N, Noguchi T, Koga T. Immunochemical and structural characterization of a serotype-specific polysaccharide antigen from *Actinobacillus actinomycetemcomitans* Y4 (serotype b). *Infect Immun* 1989; **57**: 2942–2946.
- Asikainen S, Lai C-H, Alaluusua S, Slots J. Distribution of *Actinobacillus actinomycetemcomitans* serotypes in periodontal health and disease. *Oral Microbiol Immunol* 1991; **6**: 115–118.
- Rantz LA, Randall E. Use of autoclaved extracts of hemolytic streptococci for serological grouping. *Stanford Med Bull* 1955; **13**: 290–291.
- Ouchterlony O. Diffusion-in-gel methods for immunological analysis. II. *Progr Allerg* 1962; **6**: 30–154.
- Sambrook J, Fritsch EF, Maniatis T. Molecular cloning: a laboratory manual, 2nd edn. Cold Spring Harbor, NY, Cold Spring Harbor Laboratory Press. 1989: B.13.
- Hendrickson DA, Krenz MM. Reagents and stains. In: Balows A, Hausler WJ, Herrmann KL, Isenberg HD, Shadomy HJ (eds) Manual of clinical microbiology, 5th edn. Washington, DC, American Society for Microbiology. 1991: 1289–1314.
- Ainamo J, Bay I. Problems and proposals for recording gingivitis and plaque. *Int Dent J* 1975; **25**: 229–235.
- Slots J, Reynolds HS. Long-wave UV-light fluorescence for identification of black-pigmented *Bacteroides* spp. *J Clin Microbiol* 1982; **16**: 1148–1151.
- Summanen P. Rapid disk and spot tests for the identification of anaerobes. In: Isenberg HD (ed) Clinical microbiology procedures handbook. Washington, DC, American Society for Microbiology. 1992: 2.5.6–2.5.7.
- Haffajee AD, Socransky SS, Dzink JL, Taubman MA, Ebersole JL, Smith DJ. Clinical, microbiological and immunological features of subjects with destructive periodontal diseases. *J Clin Periodontol* 1988; **15**: 240–246.
- Zambon JJ, Umemoto T, De Nardin E, Nakazawa F, Christersson FA, Genco RJ. *Actinobacillus actinomycetemcomitans* in the pathogenesis of human periodontal disease. *Adv Dent Res* 1988; **2**: 269–274.
- Dahlen G, Pipattanogovit P, Rosling B, Möller ÅJR. A comparison of two transport media for saliva and subgingival samples. *Oral Microbiol Immunol* 1993; **8**: 375–382.